

THE EXTENSION OF BOUNDED HOLOMORPHIC FUNCTIONS FROM HYPERSURFACES IN A POLYCYLINDER

by Aldo Andreotti and Wilhelm Stoll¹

Let S be an analytic subset of pure codimension 1 of a complex manifold G . Let f be a bounded, holomorphic function on S . Then, the following extension problem can be asked: *Does there exist a bounded holomorphic function F on G such that $F|_S = f$?* Of course, in general such an extension is impossible. Even in simple cases, the answer is difficult, for instance in the case of the unit disc. Alexander [1] solved the problem for the polydisc under very restrictive assumptions on S^2 . Stout [6] solved the second Cousin problem with bounds for the polydisc. His paper was the starting point for these investigations. Grauert and Lieb [2] and Ramírez de Arellano [4] solved the first Cousin problem with bounds if G is a strongly pseudoconvex open, bounded subset of \mathbb{C} with C^∞ -boundary.

Here, the problem will be solved under suitable assumptions, if G is a polycylinder, i.e., $G = G_1 \times \cdots \times G_n$, where each G_v is an open, connected bounded subset of the complex plane \mathbb{C} . As Alexander has shown by a counterexample, the problem in general is unsolvable, i.e., it is unreasonably posed. Precisely, the result is as follows: Let h be a bounded, holomorphic function on G and suppose that $S = h^{-1}(0)$ is non empty and different from $G = G_1 \times \cdots \times G_n$. A holomorphic function $f: S \rightarrow \mathbb{C}$ is said to be *strictly bounded*, if and only if there exists a finite open covering $\{U_i\}_{i \in I}$ of \bar{G} , a bounded holomorphic function f_i on $U_i \cap G$ for each $i \in I$, and a bounded holomorphic function a_{ij} on $U_i \cap U_j \cap G$ for each $(i, j) \in I \times I$ with $U_i \cap U_j \neq \emptyset$ such that

$$f|_{U_i \cap S} = f_i|_{U_i \cap S} \quad (i \in I)$$

$$f_i - f_j = a_{ij}h \quad \text{on } U_i \cap U_j \cap G$$

if $U_i \cap U_j \neq \emptyset$. Theorem 2.8 asserts that a bounded holomorphic function $F: G \rightarrow \mathbb{C}$ with $F|_S = f$ exists, if f is strictly bounded.

The proof uses a type of Grothendieck Lemma with bounds on $G_1 \times \cdots \times G_n$. The exact statement is given in Theorem 2.8.

1. The Grothendieck Lemma with Bounds

Notations. If $f: M \rightarrow \mathbf{C}$ is a function, define $\|f\|_M = \sup |f(M)|$. If M is a real or complex manifold, let $C^k(M)$ be the set of functions $f: M \rightarrow \mathbf{C}$ of class C^k , where $0 \leq k \leq \infty$ implies the usual meaning and $k = h$ means holomorphic. If $M \subseteq M_1 \times \cdots \times M_n$, let $C^k(M, k_1 \cdots k_n)$ be the set of all $f \in C^k(M)$, such that f is of class C^{k_v} in the v th variable.

If M is a set, define $M^n = M \times \cdots \times M$ (n -times). Let \mathbf{N}_0 be the set of non negative integers. For $\mu \in \mathbf{N}_0^n$, define the differential operator

$$D_\mu = D_\mu^{\bar{z}} = \frac{\partial^{\mu_1 + \cdots + \mu_n}}{\partial \bar{z}_1^{\mu_1} \cdots \partial \bar{z}_n^{\mu_n}}.$$

If M is open in \mathbf{C}^n , if $f \in C^k(M)$ and if $0 \leq p \leq k$ and $0 \leq \lambda \leq n$, let $\|f\|_M^{(p, \lambda)}$ be the maximum of all $\|D_\mu f\|_M$ where $0 \leq \mu_1 + \cdots + \mu_\lambda \leq p$ and $\mu_{\lambda+1} = \cdots = \mu_n = 0$. Define $\|f\|_M^{(p)} = \|f\|_M^{(p, n)}$. Then

$$\|f\|_M^{(0)} = \|f\|_M = \|f\|_M^{(0, \lambda)}.$$

Define

$$C_{p, \lambda}^k(M) = \{f \in C^k(M) \mid \|f\|_M^{(p, \lambda)} < \infty\},$$

$$C_p^k(M) = C_{p, n}^k(M).$$

If $K \subseteq H \subset \mathbf{C}$, let

$$\beta(K, H) = \inf \{|z - w| \mid z \in K, w \in \mathbf{C} - H\}$$

be the boundary distance from K to H . If K is compact and $H \neq \mathbf{C}$, then $\infty > \beta(K, H) > 0$. If H is a bounded subset of \mathbf{C} , its diameter is given by $\text{diam}(H) = \sup \{|z - w| \mid (z, w) \in H \times H\}$. If H is a measurable subset of \mathbf{C} , let $m(H)$ be its Lebesgue measure.

Lemma 1.1. Let M_1 be an open subset of \mathbf{C}^n with $n \geq 0$. Let M_2 be a complex manifold. For every subset H of \mathbf{C} define $H' = M_1 \times H \times M_2$. Let G be a non empty, bounded, open subset of \mathbf{C} with diameter Δ . Define $M = G'$.

Then for every open set with $\emptyset \neq H \subset \bar{H} \subset G$, a linear map

$$\kappa = \kappa_H: C^\infty(M; \infty, \infty, h) \rightarrow C^\infty(\mathbf{C}^1; \infty, \infty, h)$$

exists such that for all non negative integers s

- a) $\|\kappa(f)\|_M^{(s,n)} \leq 2\Delta \|f\|_M^{(s,n)} \leq \infty$,
 b) $\|\kappa(f)_z\|_{\tilde{C}}^{(s,n)} \leq \|f\|_M^{(s,n)} \leq \infty$,
 c) $\kappa(f)_z = f$ on H' ,
 d) if K is compact, if H and \bar{H} are open with

$$\emptyset \neq K \subset H \subset \bar{H} \subset \tilde{H} \subset \bar{\tilde{H}} \subset G,$$

then

$$\|\kappa_H(f) - \kappa_{\tilde{H}}(f)\|_{K'}^{(s,n)} \leq \frac{m(G-H)}{\beta(K, \bar{H})} \|f\|_M^{(s,n)}.$$

e) If \hat{M}_v is an open, non empty subset of M_v , if H is open with $\emptyset \neq H \subset \bar{H} \subset G$ and if r are the natural restriction maps, then the following diagram is commutative:

$$\begin{array}{ccc} C^\infty(M_1 \times G \times M_2; \infty, \infty, h) & \xrightarrow{\kappa_H} & C^\infty(M_1 \times \mathbf{C} \times M_2; \infty, \infty, h) \\ r \downarrow & & \downarrow r \\ C^\infty(\hat{M}_1 \times G \times \hat{M}_2; \infty, \infty, h) & \xrightarrow{\kappa_H} & C^\infty(\hat{M}_1 \times \mathbf{C} \times \hat{M}_2; \infty, \infty, h). \end{array}$$

Proof. Let H be open and bounded with $\bar{H} \subset G$. Take a function $\rho = \rho_H$ of class C^∞ on \mathbf{C} with compact support in G such that $0 \leq \rho \leq 1$ and such that $\rho = 1$ in a neighborhood of \bar{H} . Take $f \in C^\infty(M; \infty, \infty, h)$. For $(x, z, w) \in \mathbf{C}' = M_1 \times \mathbf{C} \times M_2$, define

$$g(x, z, w) = \begin{cases} f(x, z, w) \rho(z) & \text{if } z \in G \\ 0 & \text{if } z \in \mathbf{C} - G. \end{cases}$$

Then $g \in C^\infty(\mathbf{C}'; \infty, \infty, h)$. For $(x, z, w) \in \mathbf{C}'$, define

$$\kappa(f) = \frac{1}{2\pi i} \int_G g(x, \zeta, w) \rho(\zeta) \frac{d\zeta \wedge d\bar{\zeta}}{\zeta - z}.$$

If U contains G and has diameter $r \leq \infty$, then

$$\kappa(f) = -\frac{1}{\pi} \int_0^r \int_0^{2\pi} g(x, te^{i\phi} + z, w) e^{-i\phi} d\phi dt,$$

on U' . If $U = G$, then $r = \Delta$. The expressions for $\kappa(f)$ show $\kappa(f) \in C^\infty(M; \infty, \infty, h)$. Obviously, κ is linear. Moreover, if $z \in G$, then

$$D_\mu^{\bar{x}} \kappa(f) = -\frac{1}{\pi} \int_0^{2\pi} \int_0^\Delta (D_\mu^{\bar{x}} g)(x, te^{i\phi} + z, w) e^{-i\phi} dt d\phi.$$

Hence

$$|D_{\mu}^{\bar{x}} \kappa(f)| \leq 2\Delta \|D_{\mu}^{\bar{x}} g\|_{C'} \leq 2\Delta \|D_{\mu}^{\bar{x}} f\|_M$$

which implies a). Obviously e) is true. As is well known, $\kappa(f)_{\bar{z}} = g$ on C' (Hörmander [3, p. 3, Theorem 1.2.2]). Therefore, b) and c) are true.

Now, the assumptions of d) are made. Take $(x, z, w) \in K' = M_1 \times K \times M_2$: then

$$\begin{aligned} & |D_{\mu}^{\pi} \kappa_H(f)(x, z, w) - D_{\mu}^{\bar{x}} \kappa_{\bar{H}}(f)(x, z, w)| \\ &= \left| \frac{1}{2\pi i} \int_G D_{\mu}^{\bar{x}} f(x, \zeta, w) (\rho_H(\zeta) - \rho_{\bar{H}}(\zeta)) \frac{d\zeta \wedge d\bar{\zeta}}{\zeta - z} \right| \\ &\leq \frac{1}{2\pi} \int_{G-H} |(D_{\mu}^{\bar{x}} f)(x, \zeta, w)| \frac{d\zeta \wedge d\bar{\zeta}}{|\zeta - z|} \\ &\leq \frac{m(G-H)}{\beta(K, H)} \|D_{\mu}^{\bar{x}} f\|_M \end{aligned}$$

which implies d).

Q.E.D.

For $1 \leq p \leq n$, let $T(p, n)$ be the set of injective and increasing maps of $\{1, 2, \dots, p\}$ into $\{1, \dots, n\}$. If $\mu \in T(p, n)$ and f_1, \dots, f_n are differentiable functions, define

$$df_{\mu} = df_{\mu(s)} \wedge \dots \wedge df_{\mu(p)}.$$

If G is open in C^n , let $A^{p,q}(G)$ be the vector space of complex forms of bidegree (p, q) and of class C^{∞} on G . Define

$$Z^{p,q}(G) = \{\omega \in A^{p,q}(G) \mid \bar{\partial}\omega = 0\}.$$

Let z_1, \dots, z_n be the coordinate functions on G . If $\omega \in A^{p,q}(G)$, then

$$\omega = \sum_{\mu \in T(p, n)} \sum_{v \in T(q, n)} \omega_{\mu v} dz_{\mu} \times d\bar{z}_v$$

where $\omega_{\mu v} \in C^{\infty}(G)$. Define

$$\begin{aligned} \|\omega\|_G^{(s, \lambda)} &= \sup \{ \|\omega_{\mu v}\|_G^{(s, \lambda)} \mid (\mu, v) \in T(p, n) \times T(q, n) \}, \\ \|\omega\|_G^{(s)} &= \|\omega\|_G^{(s, n)}, \\ \|\omega\|_G &= \|\omega\|_G^{(0)} = \|\omega\|_G^{(0, \lambda)}. \end{aligned}$$

For $q \leq \lambda \leq n$, let $A_{\lambda}^{p,q}(G)$ be the set of $\omega \in A^{p,q}(G)$ with

$$\omega = \sum_{\mu \in T(p, n)} \sum_{v \in T(q, \lambda)} \omega_{\mu v} dz_{\mu} \wedge d\bar{z}_v.$$

Define $Z_\lambda^{p,q}(G) = A_\lambda^{p,q}(G) \cap Z^{p,q}(G)$. Now Lemma 1.1 extends:

Lemma 1.2. Suppose $1 \leq q \leq n$. Let M_1 be open in \mathbf{C}^{q-1} and M_2 be open in \mathbf{C}^{n-q} . If $H \subseteq \mathbf{C}$, define $H' = M_1 \times H \times M_2$. Let G be a non empty, bounded, open subset of \mathbf{C} with diameter Δ . Define $M = G'$. Then, for every open set H with $\emptyset \neq H \subset \bar{H} \subset G$, a linear map

$$K = K_H: Z_q^{p,q}(M) \rightarrow A_{q-1}^{p,q-1}(\mathbf{C}')$$

exists such that $K_H(\omega)$ is holomorphic in the variables (z_{q+1}, \dots, z_n) on M_2 if $\omega \in Z_q^{p,q}(M)$ and such that

$$\text{a) } \|K(\omega)\|_M \leq 2\Delta \|\omega\|_M,$$

$$\text{b) } \bar{\partial}K(\omega) = \omega \quad \text{on } H',$$

c) if C is compact, if H and \tilde{H} are open with

$$\emptyset \neq C \subset H \subset \bar{H} \subset \tilde{H} \subset \bar{\tilde{H}} \subset G$$

then

$$\|K_H(\omega) - K_{\tilde{H}}(\omega)\|_{C'} \leq \frac{m(G-H)}{\beta(C, H)} \|\omega\|_M,$$

d) if \hat{M}_v is an open, non empty subset of M_v for $v = 1, 2$, if H is open with $\emptyset \neq H \subset \bar{H} \subset G$ and if r denotes the natural restriction maps, then the following diagram is commutative:

$$\begin{array}{ccc} Z_q^{p,q}(M_1 \times G \times M_2) & \xrightarrow{K_H} & A_{q-1}^{p,q-1}(M_1 \times \mathbf{C} \times M_2) \\ r \downarrow & & \downarrow r \\ Z_q^{p,q}(\hat{M}_1 \times G \times \hat{M}_2) & \xrightarrow{K_H} & A_{q-1}^{p,q-1}(\hat{M}_1 \times \mathbf{C} \times \hat{M}_2). \end{array}$$

Proof. If $\omega \in Z_q^{p,q}(M)$, then

$$\omega = \sum_{\mu \in T(p,n)} \omega_\mu dz_\mu \wedge d\bar{z}_1 \wedge \dots \wedge d\bar{z}_q$$

where $\omega_{\mu\bar{z}_\rho} = 0$ for $\rho = q+1, \dots, n$. Hence $\omega_\mu \in C^\infty(M; \infty, \infty, h)$. Define the linear map K by

$$K(\omega) = (-1)^{p+q-1} \sum_{\mu \in T(p,n)} \kappa_H(\omega_\mu) dz_\mu \wedge d\bar{z}_1 \wedge \dots \wedge d\bar{z}_{q-1}.$$

Then $K_H(\omega) = K(\omega) \in A_{q-1}^{p,q-1}(\mathbf{C}')$ is holomorphic in the variables $(z_{q+1}, \dots, z_n) \in M_2$. Lemma 1.1 implies a), c) and d) immediately. Moreover,

$$\begin{aligned}
\bar{\partial}K(\omega) &= (-1)^{p+q-1} \sum_{\rho=1}^n \sum_{\mu \in T(p,n)} \kappa(\omega_\mu)_{\bar{z}_\rho} d\bar{z}_\rho \wedge dz_\mu \wedge d\bar{z}_1 \wedge \cdots \wedge d\bar{z}_{q-1} \\
&= \sum_{\mu \in T(p,n)} \kappa(\omega_\mu)_{\bar{z}_q} dz_\mu \wedge d\bar{z}_1 \wedge \cdots \wedge d\bar{z}_q \\
&= \omega
\end{aligned}$$

on H' , because $\kappa(\omega_\mu)$ is holomorphic in (z_{q+1}, \dots, z_n) . Q.E.D.

In the next lemma, the usual induction procedure will be used to prove Grothendieck's Lemma with bounds:

Lemma 1.3. *Let $1 \leq q \leq \lambda \leq n$. Suppose that $G = G_1 \times \cdots \times G_n$ where each G_v is a non empty, open subset of \mathbb{C} . Assume, that $\Delta \geq 1$ exists with $\Delta \geq \text{diam } G_v$ for $v = q, \dots, n$. Define*

$$G^\mu = G_1 \times \cdots \times G_\mu \quad G(\mu) = G_{\mu+1} \times \cdots \times G_n.$$

If $H_\mu \subseteq \mathbb{C}$ for $\mu = q, \dots, \lambda$, define

$$H^\mu = G^{q-1} \times H_q \times \cdots \times H_\mu \quad H = H^\lambda.$$

Then, for every open set $H_q \times \cdots \times H_\lambda$ with $\emptyset \neq H_\mu \subset \bar{H}_\mu \subset G_\mu$ for $\mu = q, \dots, \lambda$, a linear map

$$K = K_H: Z_\lambda^{p,q}(G) \rightarrow A_{\lambda-1}^{p,q-1}(H \times G(\lambda))$$

exists such that $K(\omega)$ is holomorphic in the variables $(z_{\lambda+1}, \dots, z_n) \in G(\lambda)$ if $\omega \in Z_\lambda^{p,q}(G)$ and such that

$$\text{a) } \|K(\omega)\|_{H \times G(\lambda)} \leq (6\Delta)^{\lambda-q+1} \|\omega\|_G^{(\lambda-q, \lambda-1)},$$

$$\text{b) } \bar{\partial}K(\omega) = \omega \quad \text{on } H \times G(\lambda),$$

c) if C_μ is compact, and if H_μ and \tilde{H}_μ are open with

$$\emptyset \neq C_\mu \subset H_\mu \subset \bar{H}_\mu \subset \tilde{H}_\mu \subset \bar{\tilde{H}}_\mu \subset G_\mu$$

or $\mu = q, \dots, \lambda$, if $C = C^\lambda \times G(\lambda)$, if $\tilde{H} = \tilde{H}^\lambda$, if $\beta = \min\{\beta(C_\mu, H_\mu) \mid \mu = q, \dots, \lambda\}$ and if $\delta = \max\{m(G_\mu - H_\mu) \mid \mu = q, \dots, \lambda\}$, then

$$\|K_H(\omega) - K_{\tilde{H}}(\omega)\|_C \leq \frac{\delta}{\beta} (24\Delta)^{\lambda-q} \|\omega\|_G^{(\lambda-q, \lambda-1)},$$

d) if \hat{G}_v is an open, non empty subset of G_v for $v = \lambda+1, \dots, n$, if H_μ is open with $\emptyset \neq H_\mu \subset \bar{H}_\mu \subset G_\mu$ for $\mu = q+1, \dots, \lambda$, then the diagram

$$\begin{array}{ccc}
Z_{\lambda}^{p,q}(G^{\lambda} \times G(\lambda)) & \xrightarrow{K_H} & A_{\lambda-1}^{p,q-1}(H \times G(\lambda)) \\
r \downarrow & & \downarrow r \\
Z_{\lambda}^{p,q}(G^{\lambda} \times \hat{G}(\lambda)) & \xrightarrow{K_H} & A_{\lambda-1}^{p,q-1}(H \times \hat{G}(\lambda))
\end{array}$$

is commutative, where r denotes the natural restriction maps.

Proof. The proof proceeds by induction for λ . Lemma 1.2 implies Lemma 1.3 for $q = \lambda$. Suppose that Lemma 1.3 is true for $\lambda - 1$ with $q < \lambda \leq n$. Then, it shall be proved for λ as follows.

Take open, non empty sets H_q, \dots, H_{λ} with $\bar{H}_{\mu} \subset G_{\mu}$. Several linear maps shall be defined:

$$\begin{aligned}
\pi: Z_{\lambda}^{p,q}(G) &\rightarrow A_{\lambda-1}^{p,q-1}(G) \\
\psi: Z_{\lambda}^{p,q}(G) &\rightarrow A_{\lambda-1}^{p,q}(G) \\
\chi: A_{\lambda-1}^{p,q-1}(G) &\rightarrow A_{\lambda-1}^{p,q-1}(G) \\
\sigma: Z_{\lambda}^{p,q}(G) &\rightarrow Z_{\lambda-1}^{p,q}(G^{\lambda-1} \times H_{\lambda} \times G(\lambda)) \\
K: Z_{\lambda}^{p,q}(G) &\rightarrow A_{\lambda-1}^{p,q-1}(H \times G(\lambda))
\end{aligned}$$

1. *Definition of π and ψ :* Take $\omega \in Z_{\lambda}^{p,q}(G)$, then $\omega = d\bar{z}_{\lambda} \wedge \alpha + \beta$ where $\alpha = \pi(\omega) \in A_{\lambda-1}^{p,q-1}(G)$ and $\beta = \psi(\omega) \in A_{\lambda-1}^{p,q}(G)$ are unique.

2. *Definition of χ .* Take $\alpha \in A_{\lambda-1}^{p,q-1}(G)$, then

$$\alpha = \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1, \lambda-1)} \alpha_{\mu v} dz_{\mu} \wedge d\bar{z}_v$$

where $\alpha_{\mu v} \in C^{\infty}(G^{\lambda-1} \times G_{\lambda} \times G(\lambda))$. Define

$$\gamma = \chi(\alpha) = \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1, \lambda-1)} \kappa_{H_{\lambda}}(\alpha_{\mu v}) dz_{\mu} \wedge d\bar{z}_v.$$

3. *Definition of σ :* If $\omega \in Z_{\lambda}^{p,q}(G)$ define

$$\sigma(\omega) = (\omega - \bar{\partial} \circ \chi \circ \pi(\omega))|_{G^{\lambda-1} \times H_{\lambda} \times G(\lambda)}.$$

Then $\bar{\partial}\sigma(\omega) = \bar{\partial}\omega = 0$. Denote $\alpha = \pi(\omega)$ and $\beta = \psi(\omega)$. Then $\omega = d\bar{z}_{\lambda} \wedge \alpha + \beta$ with

$$\begin{aligned}
\alpha &= \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1, \lambda-1)} \alpha_{\mu v} dz_{\mu} \wedge d\bar{z}_v \\
\beta &= \sum_{\mu \in T(p,n)} \sum_{v \in T(q, \lambda-1)} \beta_{\mu v} dz_{\mu} \wedge d\bar{z}_v
\end{aligned}$$

$$\begin{aligned}\bar{\partial}\omega &= -d\bar{z}_\lambda \wedge \sum_{\rho=1}^n \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1, \lambda-1)} \alpha_{\mu v \bar{z}_\rho} d\bar{z}_\rho \wedge dz_\mu \wedge d\bar{z}_v \\ &+ \sum_{\rho=1}^n d\bar{z}_\rho \wedge \sum_{\mu \in T(p,n)} \sum_{v \in T(q, \lambda-1)} \beta_{\mu v \bar{z}_\rho} dz_\mu \wedge d\bar{z}_v.\end{aligned}$$

Therefore, $\alpha_{\mu v \bar{z}_\rho} = 0$ and $\beta_{\mu v \bar{z}_\rho} = 0$ if $\rho = \lambda + 1, \dots, n$. Hence $\alpha_{\mu v}$ and $\beta_{\mu v}$ are holomorphic in the variables $(z_{\lambda+1}, \dots, z_n)$ on $G(\lambda)$, which implies the same property for $\kappa_{H_\lambda}(\alpha_{\mu v})$. Set $\gamma = \chi(\alpha)$. Then

$$\begin{aligned}\bar{\partial}\gamma &= \sum_{\rho=1}^n \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1, \lambda-1)} \kappa_{H_\lambda}(\alpha_{\mu v})_{\bar{z}_\rho} d\bar{z}_\rho \wedge dz_\mu \wedge d\bar{z}_v \\ &= d\bar{z}_\lambda \wedge \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1, \lambda-1)} \alpha_{\mu v} dz_\mu \wedge d\bar{z}_v \\ &+ \sum_{\rho=1}^{\lambda-1} \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1, \lambda-1)} \kappa_{H_\lambda}(\alpha_{\mu v})_{\bar{z}_\rho} d\bar{z}_\rho \wedge dz_\mu \wedge d\bar{z}_v \\ &= d\bar{z}_\lambda \wedge \alpha + \eta \quad \text{on } G^{\lambda-1} \times H_\lambda \times G(\lambda),\end{aligned}$$

where $\eta \in A_{\lambda-1}^{p,q}(G)$. Hence

$$\sigma(\omega) = \omega - \bar{\partial}\gamma = \beta - \eta \in Z_{\lambda-1}^{p,q}(G^{\lambda-1} \times H_\lambda \times G(\lambda)).$$

4. *Definition of $K = K_H$:* Now, let τ be the operator K which exists by induction for $G_1, \dots, G_{\lambda-1}$, H_λ , $G_{\lambda+1}, \dots, G_n$ as fixed domains and $H_q \subset \bar{H}_q \subset G_q, \dots, H_{\lambda-1} \subset \bar{H}_{\lambda-1} \subset G_{\lambda-1}$ as subdomains. Observe that $H^{\lambda-1} \times H_\lambda \times G(\lambda) = H \times G(\lambda)$. Hence

$$\tau: Z_{\lambda-1}^{p,q}(G^{\lambda-1} \times H_\lambda \times G(\lambda)) \rightarrow A_{\lambda-2}^{p,q-1}(H \times G(\lambda))$$

where $A_{\lambda-2}^{p,q-1}(H \times G(\lambda))$ is contained in $A_{\lambda-1}^{p,q-1}(H \times G(\lambda))$. Define

$$K = K_H = \chi \circ \pi + \tau \circ \sigma: Z_\lambda^{p,q}(G) \rightarrow A_{\lambda-1}^{p,q-1}(H \times G(\lambda)).$$

Take $\omega \in Z_\lambda^{p,q}(G)$. Define $\alpha = \pi(\omega)$ and $\beta = \psi(\omega)$. Then $\omega = d\bar{z}_\lambda \wedge \alpha + \beta$. Then $\gamma = \chi(\alpha) \in A_{\lambda-1}^{p,q-1}(G)$ is holomorphic in $(z_{\lambda+1}, \dots, z_n)$ on $G(\lambda)$ as shown above. Moreover, $\tau(\sigma(\omega))$ is holomorphic in $(z_{\lambda+1}, \dots, z_n)$ on $G(\lambda)$ by induction. Hence $K(\omega)$ is holomorphic in $(z_{\lambda+1}, \dots, z_n)$ on $G(\lambda)$. Moreover, $\sigma(\omega) = \omega - \bar{\partial}\gamma$ and

$$\begin{aligned}\bar{\partial}K(\omega) &= \bar{\partial}\gamma + \bar{\partial}\tau(\omega - \bar{\partial}\gamma) \\ &= \bar{\partial}\gamma + \omega - \bar{\partial}\gamma = \omega\end{aligned}$$

on $H^{\lambda-1} \times H_\lambda \times G(\lambda) = H \times G(\lambda)$ which proves b).

Now a) shall be proved: Take $\omega \in Z_{\lambda}^{p,q}(G)$ and set again $\alpha = \pi(\omega)$, $\beta = \psi(\omega)$ and $\gamma = \chi(\alpha)$. Then $\omega = d\bar{z}_{\lambda} \wedge \alpha + \beta$ and $\sigma(\omega) = \omega - \bar{\partial}\gamma$ and

$$K(\omega) = \gamma + \tau(\sigma(\omega)).$$

Obviously,

$$\begin{aligned} \|\alpha\|_G^{(s,\rho)} &\leq \|\omega\|_G^{(s,\rho)}, \\ \|\beta\|_G^{(s,p)} &\leq \|\omega\|_G^{(s,p)}, \\ \|\gamma\|_G^{(s,\lambda-1)} &\leq 2\Delta \|\alpha\|_G^{(s,\lambda-1)} \leq 2\Delta \|\omega\|_G^{(s,\lambda-1)}, \end{aligned}$$

by Lemma 1.1. Define

$$\eta = \sum_{\rho=1}^{\lambda-1} \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1,\lambda-1)} \kappa_{H_{\lambda}}(\alpha_{\mu\nu})_{\bar{z}_{\rho}} d\bar{z}_{\rho} \wedge dz_{\mu} \wedge d\bar{z}_{\nu}.$$

Then Lemma 1.1 a) implies

$$\|\eta\|_G^{(s,\lambda-1)} \leq 2\Delta \|\alpha\|_G^{(s+1,\lambda-1)} \leq 2\Delta \|\omega\|_G^{(s+1,\lambda-1)}.$$

Now, $\sigma(\omega) = \omega - \bar{\partial}\gamma = \beta - \eta$ on $\tilde{G} = G^{\lambda-1} \times H_{\lambda} \times G(\lambda)$ implies

$$\begin{aligned} \|\sigma(\omega)\|_G^{(s,\lambda-1)} &\leq \|\beta\|_G^{(s,\lambda-1)} + \|\eta\|_G^{(s,\lambda-1)} \\ &\leq \|\omega\|_G^{(s,\lambda-1)} + 2\Delta \|\omega\|_G^{(s+1,\lambda-1)} \\ &\leq 3\Delta \|\omega\|_G^{(s+1,\lambda-1)} \end{aligned}$$

for any non negative integer s . By induction

$$\begin{aligned} \|\tau(\sigma(\omega))\|_{H \times G(\lambda)} &\leq (6\Delta)^{\lambda-q} \|\sigma(\omega)\|_G^{(\lambda-q-1,\lambda+2)} \\ &\leq \frac{1}{2}(6\Delta)^{\lambda-q+1} \|\omega\|_G^{(\lambda-q,\lambda-1)}. \end{aligned}$$

Hence

$$\begin{aligned} \|K(\omega)\|_{H \times G(\lambda)} &\leq \|\gamma\|_G + \|\tau(\sigma(\omega))\|_{H \times G(\lambda)} \\ &\leq 2\Delta \|\omega\|_G + \frac{1}{2}(6\Delta)^{\lambda-q+1} \|\omega\|_G^{(\lambda-q,\lambda-1)} \\ &\leq (6\Delta)^{\lambda-q+1} \|\omega\|_G^{(\lambda-q,\lambda-1)} \end{aligned}$$

which proves a).

Now c) shall be proved: Suppose that the assumptions of c) are made. The constructions for $\tilde{H}_q, \dots, \tilde{H}_{\lambda}$ are distinguished by \sim . Then $\tilde{\pi} = \pi$ and $\tilde{\psi} = \psi$. Take $\omega \in Z_{\lambda}^{p,q}(G)$. Then

$$K_H(\omega) - K_{\tilde{H}}(\omega) =$$

$$\chi(\pi(\omega)) - \tilde{\chi}(\pi(\omega)) + \tau(\sigma(\omega) - \tilde{\sigma}(\omega)) + \tau(\tilde{\sigma}(\omega)) - \tilde{\tau}(\tilde{\sigma}(\omega)).$$

Define $\alpha = \pi(\omega) = \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1, \lambda-1)} \alpha_{\mu v} dz_\mu \wedge d\bar{z}_v$. Then

$$\chi(\alpha) - \tilde{\chi}(\alpha) = \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1, \lambda-1)} (\kappa_{H_\lambda}(\alpha_{\mu v}) - \kappa_{\tilde{H}_\lambda}(\alpha_{\mu v})) dz_\mu \wedge d\bar{z}_v$$

where κ_{H_λ} and $\kappa_{\tilde{H}_\lambda}$ are formed for $G^{\lambda-1} \times H_\lambda \times G(\lambda)$ and $G^{\lambda-1} \times \tilde{H}_\lambda \times G(\lambda)$ respectively. Therefore Lemma 1.1 implies

$$\|\chi(\alpha) - \tilde{\chi}(\alpha)\|_C \leq \frac{m(G_\lambda - H_\lambda)}{\beta(C_\lambda, H_\lambda)} \|\alpha\|_G \leq \frac{\delta}{\beta} \|\omega\|_G.$$

Now

$$\tau = K_{H^{\lambda-1}}: Z_{\lambda-1}^{p,q}(G^{\lambda-1} \times H_\lambda \times G(\lambda)) \rightarrow A_{\lambda-1}^{p,q}(H^{\lambda-1} \times H_\lambda \times G(\lambda))$$

$$\tilde{\tau} = K_{\tilde{H}^{\lambda-1}}: Z_{\lambda-1}^{p,q}(G^{\lambda-1} \times \tilde{H}_\lambda \times G(\lambda)) \rightarrow A_{\lambda-1}^{p,q}(\tilde{H}^{\lambda-1} \times \tilde{H}_\lambda \times G(\lambda)).$$

Define

$$\tilde{\tau}_0 = K_{\tilde{H}^{\lambda-1}}: Z_{\lambda-1}^{p,q}(G^{\lambda-1} \times H_\lambda \times G(\lambda)) \rightarrow A_{\lambda-1}^{p,q}(\tilde{H}^{\lambda-1} \times H_\lambda \times G(\lambda)).$$

The induction assumption d) implies $\tilde{\tau}(\tilde{\sigma}(\omega)) = \tilde{\tau}_0(\tilde{\sigma}(\omega))$. The induction assumption c) implies

$$\begin{aligned} & \|\tilde{\tau}(\tilde{\sigma}(\omega)) - \tau(\tilde{\sigma}(\omega))\|_C \\ & \leq \|\tilde{\tau}_0(\tilde{\sigma}(\omega)) - \tau(\tilde{\sigma}(\omega))\|_{C^{\lambda-1} \times H_\lambda \times G(\lambda)} \\ & \leq \frac{\delta}{\beta} (24\Delta)^{\lambda-q-1} \|\tilde{\sigma}(\omega)\|_{G^{\lambda-1} \times \tilde{H}_\lambda \times G(\lambda)}^{(\lambda-q-1, \lambda-2)} \\ & \leq \frac{\delta}{\beta} (24\Delta)^{\lambda-q-1} \|\tilde{\sigma}(\omega)\|_{G^{\lambda-1} \times \tilde{H}_\lambda \times G(\lambda)}^{(\lambda-q-1, \lambda-1)} \\ & \leq \frac{\delta}{4\beta} (24\Delta)^{\lambda-q} \|\omega\|_G^{(\lambda-q, \lambda-1)}. \end{aligned}$$

Now, $\sigma(\omega) = \beta - \eta$ on $G^{\lambda-1} \times H_\lambda \times G(\lambda)$ and $\tilde{\sigma}(\omega) = \beta - \tilde{\eta}$ on $G^{\lambda-1} \times \tilde{H}_\lambda \times G(\lambda)$ imply

$$\sigma(\omega) - \tilde{\sigma}(\omega) = \eta - \tilde{\eta} =$$

$$\sum_{\rho=1}^{\lambda-1} \sum_{\mu \in T(p,n)} \sum_{v \in T(q-1, \lambda-1)} (\kappa_{H_\lambda}(\alpha_{\mu v})_{\bar{z}_\rho} - \kappa_{\tilde{H}_\lambda}(\alpha_{\mu v})_{\bar{z}_\rho}) d\bar{z}_\rho \wedge dz_\mu \wedge d\bar{z}_v$$

on $G^{\lambda-1} \times H_\lambda \times G(\lambda)$ where $\alpha_{\mu\nu} \in C^\infty(G^{\lambda-1} \times G_\lambda \times G(\lambda); \infty, \infty, h)$. Let U be an open neighborhood of C_λ with $\bar{U} \subset H_\lambda$ such that $\beta(C_\lambda, H_\lambda) \leq 2\beta(\bar{U}, H_\lambda)$. Lemma 1.1 d) implies

$$\begin{aligned} & \| \kappa_{H_\lambda}(\alpha_{\mu\nu}) - \kappa_{\tilde{H}_\lambda}(\alpha_{\mu\nu}) \|_{G^{\lambda-1} \times U \times G(\lambda)}^{(s, \lambda-1)} \\ & \leq \frac{m(G_\lambda - H_\lambda)}{\beta(\bar{U}, H_\lambda)} \| \alpha_{\mu\nu} \|_{G^{\lambda-1} \times G_\lambda \times G(\lambda)}^{(s, \lambda-1)} \\ & \leq \frac{2\delta}{\beta} \| \omega \|_G^{(s, \lambda-1)}. \end{aligned}$$

Hence

$$\| \kappa_{H_\lambda}(\alpha_{\mu\nu})_{\bar{z}_\rho} - \kappa_{\tilde{H}_\lambda}(\alpha_{\mu\nu})_{\bar{z}_\rho} \|_{G^{\lambda-1} \times U \times G(\lambda)}^{(s, \lambda-1)} \leq \frac{2\delta}{\beta} \| \omega \|_G^{(s+1, \lambda-1)}$$

for $\rho = 1, \dots, \lambda-1$, which implies

$$\| \sigma(\omega) - \tilde{\sigma}(\omega) \|_{G^{\lambda-1} \times U \times G(\lambda)}^{(s, \lambda-1)} \leq \frac{2\delta}{\beta} \| \omega \|_G^{(s+1, \lambda-1)}.$$

Define

$$\tau_0 = K_{H^{\lambda-1}}: Z_{\lambda-1}^{p, q}(G^{\lambda-1} \times U \times G(\lambda)) \rightarrow A_{\lambda-1}^{p, q-1}(H^{\lambda-1} \times U \times G(\lambda)).$$

The induction assumption d) implies $\tau_0(\sigma(\omega) - \tilde{\sigma}(\omega)) = \tau(\sigma(\omega) - \tilde{\sigma}(\omega))$.

The induction assumption a) implies

$$\begin{aligned} & \| \tau(\sigma(\omega) - \tilde{\sigma}(\omega)) \|_C \\ & \leq \| \tau_0(\sigma(\omega) - \tilde{\sigma}(\omega)) \|_{H^{\lambda-1} \times U \times G(\lambda)} \\ & \leq (6\Delta)^{\lambda-q} \| \sigma(\omega) - \tilde{\sigma}(\omega) \|_{G^{\lambda-1} \times U \times G(\lambda)}^{(\lambda-q-1, \lambda-1)} \\ & \leq \frac{\delta}{2\beta} (24\Delta)^{\lambda-q} \| \omega \|_G^{(\lambda-q, \lambda-1)}. \end{aligned}$$

Therefore

$$\begin{aligned} & \| K_H(\omega) - K_{\tilde{H}}(\omega) \|_C \\ & \leq \| \chi(\alpha) - \tilde{\chi}(\alpha) \|_C + \| \tau(\sigma(\omega) - \tilde{\sigma}(\omega)) \|_C + \| \tau(\tilde{\sigma}(\omega)) - \tilde{\tau}(\tilde{\sigma}(\omega)) \|_C \\ & \leq \frac{\delta}{\beta} \| \omega \|_G + \frac{\delta}{2\beta} (24\Delta)^{\lambda-q} \| \omega \|_G^{(\lambda-q, \lambda-1)} + \frac{\delta}{4\beta} (24\Delta)^{\lambda-q} \| \omega \|_G^{(\lambda-q, \lambda-1)} \\ & \leq \frac{\delta}{\beta} (24\Delta)^{\lambda-q} \| \omega \|_G^{(\lambda-q, \lambda-1)} \end{aligned}$$

which proves c).

Now d) shall be proved: Suppose that the assumptions of d) are made. The constructions for $\hat{G}_{\lambda+1}, \dots, \hat{G}_n$ are distinguished by \wedge . Then $\hat{\pi} = \pi$ and $\hat{\psi} = \psi$. Take $\omega \in Z_{\lambda}^{p,q}(\hat{G})$. Then $\pi(\omega) = \hat{\pi}(\omega)$ and $\psi(\omega) = \hat{\psi}(\omega)$. Lemma 1.1 c) implies $\chi(\pi(\omega)) = \hat{\chi}(\pi(\omega)) = \hat{\chi}(\hat{\pi}(\omega))$. Hence

$$\hat{\sigma}(\omega) = \omega - \bar{\partial}\hat{\chi}(\hat{\pi}(\omega)) = \omega - \bar{\partial}\chi(\pi(\omega)) = \sigma(\omega).$$

Induction assumption d) implies $\hat{\tau}(\hat{\sigma}(\omega)) = \tau(\hat{\sigma}(\omega)) = \tau(\sigma(\omega))$. Therefore

$$\hat{K}_H(\omega) = \hat{\chi}(\hat{\pi}(\omega)) + \hat{\tau}(\hat{\sigma}(\omega)) = \chi(\pi(\omega)) + \tau(\sigma(\omega)) = K_H(\omega),$$

which proves d). Q.E.D.

For $\lambda = n$, the following theorem is obtained.

Theorem 1.4. Let $1 \leq q \leq n$. Suppose that $G = G_1 \times \dots \times G_n$, where each G_v is a non empty, open subset of \mathbb{C} . Assume that $\Delta \geq 1$ exists such that $\text{diam } G_v \leq \Delta$ for $v = q, \dots, n$. Then, for every open set

$$H = G_1 \times \dots \times G_{q-1} \times H_q \times \dots \times H_n \neq \emptyset$$

with $\bar{H}_v \subset G_v$ for $v = q, \dots, n$ a linear map

$$K_H: Z^{p,q}(G) \rightarrow A^{p,q-1}(H)$$

exists such that

$$\text{a) } \|K_H(\omega)\|_H \leq (6\Delta)^{n-q-1} \|\omega\|_G^{(n-q)}$$

$$\text{b) } \bar{\partial}K_H(\omega) = \omega \quad \text{on } H,$$

$$\text{c) if } C_\mu \text{ is compact, if } H_\mu \text{ and } \bar{H}_\mu \text{ are open and if}$$

$$\emptyset \neq C_\mu \subset H_\mu \subset \bar{H}_\mu \subset \bar{H}_\mu \subset \bar{H}_\mu \subset G_\mu$$

for $\mu = q, \dots, n$, if

$$C = G_1 \times \dots \times G_{q-1} \times C_q \times \dots \times C_n$$

$$\bar{H} = G_1 \times \dots \times G_{q-1} \times \bar{H}_q \times \dots \times \bar{H}_n$$

$$H = G_1 \times \dots \times G_{q-1} \times H_q \times \dots \times H_n$$

$$\beta = \min\{\beta(C_\mu, H_\mu) \mid \mu = q, \dots, n\}$$

$$\delta = \max\{m(G_\mu - H_\mu) \mid \mu = q, \dots, n\}$$

then

$$\|K_H(\omega) - K_{\bar{H}}(\omega)\|_C \leq \frac{\delta}{\beta} (24\Delta)^{n-q} \|\omega\|_G^{(n-q)}.$$

Let G be an open subset of \mathbf{C}^n . Let $\tilde{A}^{p,q}(G)$ be the vector space of continuous forms of bidegree (p, q) on G . Take $\omega \in Z^{p,q}(G)$ with $q \geq 1$, then $\psi \in \tilde{A}^{p,q-1}(G)$ is said to be a *strong solution* of the $\bar{\partial}$ -problem of ω on G , if and only if a sequence $\{\psi_v\}_{v \in \mathbf{N}}$ of forms $\psi_v \in A^{p,q-1}(G)$ exists such that the following conditions hold:

1. For every compact subset C of G there exists a number $v_C \in \mathbf{N}$ such that $\bar{\partial}\psi_v = \omega$ on C for all $v \geq v_C$.

2. For every compact subset C of G

$$\|\psi - \psi_v\|_C \rightarrow 0 \quad \text{for } v \rightarrow \infty.$$

If this is so write $\omega = \bar{\partial}! \psi$. Define the vector spaces

$$Z^{p,q}(G; s) = \{\omega \in Z^{p,q}(G) \mid \|\omega\|_G^{(s)} < \infty\}$$

$$\tilde{A}^{p,q}(G) = \{\omega \in \tilde{A}^{p,q}(G) \mid \|\omega\|_G < \infty\}$$

$$A^{p,q}(G; s) = \{\omega \in A^{p,q}(G) \mid \|\omega\|_G^{(s)} < \infty\}.$$

Theorem 1.5. Let $1 \leq q \leq n$. Suppose that G_v is an open, non empty subset of \mathbf{C} for $v = 1, \dots, n$. Assume that G_q, \dots, G_n are bounded. Define $G = G_1 \times \dots \times G_n$. Then a linear map

$$L: Z^{p,q}(G; n-q) \rightarrow \tilde{A}^{p,q-1}(G)$$

exists such that $\bar{\partial}! L(\omega) = \omega$ for all $\omega \in Z^{p,q}(G; n-q)$. If $\Delta \geq 1$ is such that $\text{diam } G_v \leq \Delta$ for $v = q, \dots, n$ and if $\omega \in Z^{p,q}(G; n-q)$, then

$$\|L(\omega)\|_G \leq (6\Delta)^{n-q+1} \|\omega\|_G^{(n-q)}.$$

Proof. For each $\mu = q, \dots, n$ choose a sequence $\{H_{\mu v}\}_{v \in \mathbf{N}}$ of open, non empty sets $H_{\mu v}$ with

$$H_{\mu v} \subset \bar{H}_{\mu v} \subset H_{\mu v+1} \subset \bar{H}_{\mu v+1} \subset G_\mu$$

such that $G_\mu = \bigcup_{v \in \mathbf{N}} H_{\mu v}$

$$H^v = G_1 \times \dots \times G_{q-1} \times H_{qv} \times \dots \times H_{nv}.$$

Theorem 1.4 gives a linear map

$$Q_v = K_{H^{v+2}}: Z^{p,q}(G, n-q) \rightarrow A^{p,q-1}(H^{v+2}).$$

For each $v \in \mathbf{N}$, take a C^∞ -function ρ_v on G with support in H^{v+1} and with $\rho_v|_{\bar{H}^v} = 1$ where $0 \leq \rho_v \leq 1$ on G . For $\omega \in Z^{p,q}(G, n-q)$ define $L_v(\omega) = \rho_v Q_v$ on H^{v+2} and $L_v(\omega) = 0$ on $G - H^{v+2}$. Then $L_v(\omega) \in A^{p,q-1}(G)$.

Obviously, the map L_v is linear. Take $\Delta \geq 1$ with $\Delta \geq \text{diam } G_v$ for $v = q, \dots, n$.

Let U be any open subset of G such that \bar{U} is compact. Then a number $v_0(U) \in N$ exists such that $H^v \supset \bar{U}$ for $v \geq v_0(U)$. Therefore

$$\|L_v(\omega)\|_U = \|Q_v(\omega)\|_U \leq (6\Delta)^{n-q+1} \|\omega\|_G^{n-q} < \infty$$

for $v \geq v_0(U)$ and $\omega \in Z^{p,q}(G; n-q)$. For $v \geq v_0(U)$ define $\delta_v = \max\{m(G_\mu - H_{\mu v+2}) \mid \mu = q, \dots, n\}$. Then $\delta_v \rightarrow 0$ for $v \rightarrow \infty$. Define

$$\beta(v) = \min\{\beta(\bar{U}, H_{\mu v+2}) \mid \mu = q, \dots, n\}.$$

Then $\beta(v) \geq \beta = \beta(v_0(U)) > 0$ for all $v \geq v_0(U)$. If $\mu > v \geq v_0(U)$, then

$$\begin{aligned} \|L_v(\omega) - L_\mu(\omega)\|_{\bar{U}} &= \|Q_v(\omega) - Q_\mu(\omega)\|_{\bar{U}} \\ &\leq \frac{\delta_v}{\beta} (24\Delta)^{n-q} \|\omega\|_G^{(n-q)}. \end{aligned}$$

Hence $\{L_v(\omega)\}_{v \in N}$ is a Cauchy sequence on G and converges uniformly on every compact subset of G . Then

$$L(\omega) = \lim_{v \rightarrow \infty} L_v(\omega) \in \tilde{A}^{p,q-1}(G).$$

The map L is linear. Moreover

$$\|L(\omega)\|_{\bar{U}} \leq (6\Delta)^{n-q+1} \|\omega\|_G^{(n-q)}$$

for every open, relative compact subset U of G . Hence

$$\|L(\omega)\|_G \leq (6\Delta)^{n-q+1} \|\omega\|_G^{(n-q)} < \infty,$$

which implies $L(\omega) \in \tilde{A}^{p,q-1}(G)$.

If C is compact in G , take an open neighborhood U of C with \bar{U} compact and $\bar{U} \subset G$. If $v \geq v_0(U)$, then

$$\tilde{\partial} L_v(\omega) = \tilde{\partial} Q_v(\omega) = \omega \text{ on } C$$

because $C \subseteq U \subseteq H^v$. Moreover,

$$\|L_v(\omega) - L(\omega)\|_{\bar{U}} \leq \frac{\delta_v}{\beta} (24\Delta)^{n-q} \|\omega\|_G^{(n-q)}.$$

Hence $\|L_v(\omega) - L(\omega)\|_C \rightarrow 0$ for $v \rightarrow \infty$. Consequently, $\tilde{\partial}! L(\omega) = \omega$.

Q.E.D.

If $q = 1$, the situation can easily be improved:

Lemma 1.6. Let $G \neq \emptyset$ be open in \mathbb{C}^n . Suppose that $\omega \in Z^{p,1}(G)$ and $\psi \in \tilde{A}^{p,0}(G)$ are given with $\omega = \bar{\partial}! \psi$. Then $\psi \in A^{p,0}(G)$ is of class C^∞ and $\omega = \bar{\partial} \psi$.

Proof. Let $\{\psi_v\}_{v \in N}$ be a sequence of forms $\psi_v \in A^{p,0}(G)$ satisfying the conditions of the definition of $\omega = \bar{\partial}! \psi$. Let U be any open, relative compact subset of G with $\bar{U} \subseteq G$ such that $\omega = \bar{\partial} \phi$ on U where $\phi \in A^{p,0}(U)$. A number $v_{\bar{u}} \in N$ exists such that $\bar{\partial} \psi_v = \omega = \bar{\partial} \phi$ on U for all $v \geq v_{\bar{u}}$. Hence $\chi_v = \psi_v - \phi$ is a holomorphic form on U because $\bar{\partial} \chi_v = 0$ if $v \geq v_{\bar{u}}$. Because $\psi_v \rightarrow \psi$ for $v \rightarrow \infty$ uniformly on U , it also follows that $\chi_v \rightarrow \chi = \psi - \phi$ for $v \rightarrow \infty$ uniformly on U . Hence χ is holomorphic on U , which implies that ψ is of class C^∞ on U and that $0 = \bar{\partial} \chi = \bar{\partial} \psi - \bar{\partial} \phi$ or $\bar{\partial} \psi = \omega$ on U . Because every point of G has such a neighborhood, $\psi \in A^{p,0}(G)$ and $\bar{\partial} \psi = \omega$ on G . Q.E.D.

Because $\bar{\partial}! L(\omega) = \omega$ Theorem 1.5 and Lemma 1.6 imply

Theorem 1.7. Assume that $G = G_1 \times \cdots \times G_n \neq \emptyset$ is open and bounded in \mathbb{C}^n with $G_v \subseteq \mathbb{C}$ for $v = 1, \dots, n$. Let $\Delta \geq 1$ be such that $\text{diam } G_v \leq \Delta$ for $v = 1, \dots, n$. Then a linear map

$$L: Z^{p,1}(G; n-1) \rightarrow A^{p,0}(G; 0)$$

exists such that $\bar{\partial} L(\omega) = \omega$ and

$$\|L(\omega)\|_G \leq (6\Delta)^n \|\omega\|^{(n-1)} < \infty.$$

Observe, that $A^{0,0}(G, 0)$ is the normed space of bounded complex valued C^∞ -functions on G .

2. Extension of Bounded Functions

Let $G \neq \emptyset$ be open in the complex manifold M . If $U \neq \emptyset$ is open in the Hausdorff space \bar{G} , let $B_G(U)$ be the Banach space of bounded holomorphic functions on $U \cap G$. Observe that U is open in \bar{G} and $U \cap G \neq \emptyset$ is open in M . If V is open in \bar{G} with $\emptyset \neq V \subseteq U$, the restriction map $r_V^U: B_G(U) \rightarrow B_G(V)$ is defined. Then $B = B_G = \{B_G(U), r_V^U\}$ is the pre-sheaf of bounded holomorphic functions on \bar{G} . It induces the sheaf $\mathfrak{B} = \mathfrak{B}(G)$ of germs of bounded holomorphic functions on \bar{G} . If \mathfrak{D} is the sheaf of germs of holomorphic functions on M , then $\mathfrak{D}|_G = \mathfrak{B}|_G$, but $\mathfrak{D}_x \neq \mathfrak{B}_x$ may hold if $x \in \bar{G} - G$; of course $\mathfrak{B}_x \supseteq \mathfrak{D}_x$ for all $x \in \bar{G}$.

If $U \neq \emptyset$ is open in the Hausdorff space \bar{G} , let $\Gamma(U, \mathfrak{B})$ be the algebra of sections of \mathfrak{B} over U . If V is open in \bar{G} with $\emptyset \neq V \subseteq U$, the restriction map $r_V^U: \Gamma(U, \mathfrak{B}) \rightarrow \Gamma(V, \mathfrak{B})$ is defined. Then $\Gamma(\mathfrak{B}) = \{\Gamma(U, \mathfrak{B}), r_V^U\}$ is the

canonical presheaf of \mathfrak{B} . Here, $\Gamma(U, \mathfrak{B})$ can be identified with an algebra of holomorphic functions on $U \cap G$, where $f \in \Gamma(U, \mathfrak{B})$ if and only if $f|_{(V \cap G)} \in B_G(V)$ for every open set V in \bar{G} whose closure \bar{V} is compact and contained in U . Observe $B_G(U) \subseteq \Gamma(U, \mathfrak{B})$ and that B_G is a subpresheaf of $\Gamma(\mathfrak{B})$ under this identification. If \bar{G} is compact, then $\Gamma(\bar{G}, \mathfrak{B}) = B_G(\bar{G})$ is the algebra of bounded holomorphic functions on G .

Let $\mathfrak{U} = \{U_i\}_{i \in I}$ be a covering of \bar{G} by non empty subsets U_i , which are open in the space \bar{G} . For $p \geq 0$, define

$$I(p) = \{(i_0, \dots, i_p) \mid U_{i_0} \cap \dots \cap U_{i_p} \neq \emptyset\}.$$

Then $I = I(0)$. For $i = (i_0, \dots, i_p) \in I(p)$ define $U_i = U_{i_0} \cap \dots \cap U_{i_p}$. Let \mathfrak{S}_{p+1} be the group of permutations on $(p+1)$ elements. If $\pi \in \mathfrak{S}_{p+1}$, define $\pi(i) = (i_{\pi(0)}, \dots, i_{\pi(p)})$ if $i = (i_0, \dots, i_p)$. Then $\pi(I(p)) = I(p)$. A p -cochain f on \mathfrak{U} in B is a family $f = \{f_i\}_{i \in I(p)}$ where $f_i \in B_G(U_i)$ and $f_{\pi(i)} = (\text{sign } \pi) \cdot f_i$ for each $i \in I(p)$. The set $C^p(\mathfrak{U}, B)$ of p -cochains forms a vector space over \mathbb{C} . A linear map $\delta: C^p(\mathfrak{U}, B) \rightarrow C^{p+1}(\mathfrak{U}, B)$ is defined by

$$(\delta f)_{i_0 \dots i_{p+1}} = \sum_{v=0}^{p+1} (-1)^v f_{i_0 \dots i_{v-1} i_{v+1} \dots i_{p+1}}$$

where each term is restricted to $G \cap U_i$ with $i = (i_0, \dots, i_p)$. Then $\delta \circ \delta = 0$. The vector space $Z^p(\mathfrak{U}, B) = \{f \in C^p(\mathfrak{U}, B) \mid \delta f = 0\}$ is the set of p -cocycles on \mathfrak{U} in B and contains $\delta C^{p-1}(\mathfrak{U}, B)$. For $p = 0$ define $H^0(\mathfrak{U}, B) = Z^0(\mathfrak{U}, B)$ and for $p \geq 1$ define

$$H^p(\mathfrak{U}, B) = Z^p(\mathfrak{U}, B) / \delta C^{p-1}(\mathfrak{U}, B)$$

as the p th cohomology group on \mathfrak{U} in B . If \mathfrak{B} is a refinement of \mathfrak{U} , then a linear map $r_{\mathfrak{B}}^{\mathfrak{U}}: H^p(\mathfrak{U}, B) \rightarrow H^p(\mathfrak{B}, B)$ is defined such that $r_{\mathfrak{B}}^{\mathfrak{B}} \circ r_{\mathfrak{B}}^{\mathfrak{U}} = r_{\mathfrak{B}}^{\mathfrak{U}}$ if \mathfrak{B} is a refinement of \mathfrak{B} . Hence the direct limit of $\{H^p(\mathfrak{U}, B), r_{\mathfrak{B}}^{\mathfrak{U}}\}$ exists and is $\{H^p(\bar{G}, B), r^{\mathfrak{U}}\}$ where $r^{\mathfrak{U}}: H^p(\mathfrak{U}, B) \rightarrow H^p(\bar{G}, B)$ is the residual map and where $H^p(\bar{G}, B)$ is the p th cohomology group in \bar{G} with coefficients in B . Observe $H^0(\mathfrak{U}, B) = H^0(\bar{G}, B) = H^0(\bar{G}, B) = \Gamma(\bar{G}, \mathfrak{B})$ with $r_{\mathfrak{B}}^{\mathfrak{U}}$ and $r^{\mathfrak{U}}$ being the identity.

If in these definitions the presheaf $B = B_G$ is replaced by the presheaf $\Gamma(\mathfrak{B})$, the vector spaces

$$C^p(\mathfrak{U}, \Gamma(\mathfrak{B})) = C^p(\mathfrak{U}, \mathfrak{B})$$

$$Z^p(\mathfrak{U}, \Gamma(\mathfrak{B})) = Z^p(\mathfrak{U}, \mathfrak{B})$$

$$H^p(\mathfrak{U}, \Gamma(\mathfrak{B})) = H^p(\mathfrak{U}, \mathfrak{B})$$

$$H^p(\bar{G}, \Gamma(\mathfrak{B})) = H^p(\bar{G}, \mathfrak{B})$$

are defined. The inclusion $B \subseteq \Gamma(\mathfrak{B})$ of presheaves defines an isomorphism $H^p(\bar{G}, B) \approx H^p(\bar{G}, \mathfrak{B})$, because \bar{G} is paracompact.

Suppose that \bar{G} is compact. Then every open covering on \bar{G} has a finite refinement. Therefore, the previous considerations can be restricted to finite, open coverings. If \mathfrak{U} is such a finite, open covering of \bar{G} , and if $f \in C^p(\mathfrak{U}, B)$ define

$$\|f\| = \max \{\|f_i\| \mid i \in I(p)\}.$$

Then $\|\cdot\|$ is a norm on $C^p(\mathfrak{U}, B)$ and $C^p(\mathfrak{U}, B)$ is a Banach space with this norm. Moreover, $\|\delta f\| \leq (p+1)\|f\|$. Hence δ is continuous, and $Z^p(\mathfrak{U}, B)$ is again a Banach space with this norm.

Theorem 2.1. *Let $G = G_1 \times \cdots \times G_n \neq \emptyset$ be open and bounded in \mathbf{C}^n with $G_v \subset \mathbf{C}$ for $v = 1, \dots, n$. Let B be the presheaf of germs of bounded holomorphic functions on \bar{G} . Let \mathfrak{U} be a finite open covering of the compact Hausdorff space \bar{G} . Then a continuous linear map*

$$\sigma: Z^1(\mathfrak{U}, B) \rightarrow C^0(\mathfrak{U}, B)$$

exists such that that $\delta \circ \sigma$ is the identity on $Z^1(\mathfrak{U}, B)$. Especially, $H^1(\mathfrak{U}, B) = 0$.

Proof. Take a partition $\{\rho_i\}_{i \in I}$ of unity by non negative C^∞ -functions ρ_i on \mathbf{C}^n , such that $\text{supp } \rho_i$ is compact and $\bar{G} \cap \text{supp } \rho_i \subseteq U_i$ for each $i \in I$ and such that $\sum_{i \in I} \rho_i = 1$ on \bar{G} . Now a continuous linear map $\alpha: Z^1(\mathfrak{U}, B) \rightarrow Z^{0,1}(G; n-1)$ will be defined.

Take a number $C > 1$ such that

$$\|\rho_{i\bar{z}_\lambda}\|_{U_i}^{(s)} \leq C$$

for all $i \in I$, all $\lambda = 1, \dots, n$, and all $s = 0, \dots, n$. Take

$$f = \{f_{ij}\}_{(i,j) \in I(1)}$$

in $Z^1(\mathfrak{U}, B)$. Then $f_{ij} = -f_{ji}$ on $U_i \cap U_j \cap G$ if $(i,j) \in I(1)$ and $f_{ij} + f_{jk} + f_{ki} = 0$ on $U_i \cap U_j \cap U_k \cap G$ if $(i,j,k) \in I(2)$. Define $f_{ij}(x) = 0$ if $x \in G - U_i \cap U_j$. Define $g_i = \sum_{k \in I} \rho_k f_{ki}$ on $U_i \cap G$. Then g_i is a function of class C^∞ on U_i with $\|g_i\|_{U_i \cap G} \leq \|f\|$. Moreover,

$$g_j - g_i = \sum_{k \in I} \rho_k (f_{kj} + f_{ik}) = \sum_{k \in I} \rho_k f_{ij} = f_{ij}$$

on $U_i \cap U_j \cap G$ if $(i,j) \in I(1)$. Because f_{ij} is holomorphic $\bar{\partial} g_j = \bar{\partial} g_i$ on $U_i \cap U_j \cap G$. Hence $\omega \in Z^{0,1}(G)$ exists uniquely, such that $\omega|_{U_i \cap G} = \bar{\partial} g_i$. Then

$$\omega = \sum_{\lambda=1}^n \omega_{\lambda} d\bar{z}_{\lambda}$$

on G where

$$\omega_{\lambda} = \sum_{k \in I} \rho_{k\bar{z}_{\lambda}} f_{ki} \text{ on } U_i.$$

If $\mu \in N_0^n$, then

$$D_{\mu}^{\bar{z}} \omega_{\lambda} = \sum_{k \in I} (D_{\mu}^{\bar{z}} \rho_{k\bar{z}_{\lambda}}) f_{ki},$$

which implies

$$\|\omega\|_G^{(n-1)} \leq \max_{i \in I} \|\omega\|_{U_i \cap G}^{(n-1)} \leq C \|f\| < \infty.$$

Define $\alpha(f) = \omega$. Then α is a linear map into $Z^{0,1}(G; n-1)$ with $\|\alpha(f)\|_G^{(n-1)} \leq C \|f\|$. Hence α is continuous.

According to Theorem 1.7, a continuous, linear map $L: Z^{0,1}(G; n-1) \rightarrow A^{0,0}(G, 0)$ with $\bar{\partial}L(\phi) = \phi$ exists, where $A^{0,0}(G, 0)$ is the normed space of bounded C^{∞} -functions on G . Define $\psi = L(\alpha(f))$ on G . Define $h_i = g_i - \psi$ on $U_i \cap G$ for each $i \in I$. Then $\bar{\partial}h_i = \bar{\partial}g_i - \bar{\partial}\psi = \omega - \omega = 0$ on $U_i \cap G$. Hence h_i is holomorphic on $U_i \cap G$. If $\Delta \geq 1$, such that $\text{diam } G_v \leq \Delta$ for $v = 1, \dots, n$. Then

$$\|h_i\|_{U_i \cap G} \leq C(6\Delta)^n \|f\| + \|f\| < \infty.$$

Hence, $h = \{h_i\}_{i \in I} \in C^0(\mathcal{U}, B)$. Define $h = \sigma(f)$. Obviously, σ is linear. Because

$$\|\sigma(f)\| \leq 2C(6\Delta)^n \|f\|$$

σ is continuous. If $(i, j) \in I(1)$, then

$$(\delta\sigma(f))_{ij} = (\delta h)_{ij} = h_j - h_i = g_j - g_i = f_{ij}.$$

Therefore, $(\delta \circ \sigma)(f) = f$.

Q.E.D.

Theorem 2.2. Let $G = G_1 \times \dots \times G_n \neq \emptyset$ be open and bounded in \mathbb{C}^n with $G_v \subset \mathbb{C}$ for $v = 1, \dots, n$. Let \mathfrak{B} be the sheaf of germs of bounded holomorphic functions on \bar{G} . Then

$$H^1(\bar{G}, \mathfrak{B}) = 0.$$

Proof. Take $h \in H^1(\bar{G}, B) \approx H^1(\bar{G}, \mathfrak{B})$. Then a finite, open covering \mathcal{U} of the compact Hausdorff space \bar{G} exists such that $f \in H^1(\mathcal{U}, B)$ with $h = r^{\mathcal{U}}(f)$ exists. By Theorem 2.1, $f = 0$, hence $h = 0$. Q.E.D.

Let $G \neq \emptyset$ be open in the complex manifold M . Let \mathfrak{B} be the sheaf of germs of bounded holomorphic functions on \bar{G} . If U is open in \bar{G} , if $x \in U$ and if $h \in B_G(U)$ (or $h \in \Gamma(U, \mathfrak{B})$), then h defines a germ $h_x \in \mathfrak{B}_x$. Define $h\mathfrak{B}|U = \bigcup_{x \in U} \{fh_x | f \in \mathfrak{B}_x\}$. Then $h\mathfrak{B}|U$ is a subsheaf of $\mathfrak{B}|U$ generated by h . A subsheaf \mathfrak{F} of \mathfrak{B} is said to be a principal ideal sheaf of \mathfrak{B} if and only if for every $x \in \bar{G}$ an open neighborhood U of x in \bar{G} and an element $h \in B_G(U)$ exist such that $\mathfrak{F}|U = h\mathfrak{B}|U$. If it is possible to choose $U = \bar{G}$, then \mathfrak{F} is called a globally principal ideal sheaf in \mathfrak{B} .

For any sheaf \mathfrak{F} on \bar{G} , and any open subset U of the space \bar{G} , denote by $\Gamma(U, \mathfrak{F})$ the set of continuous sections in \mathfrak{F} over U . If \mathfrak{F} is a subsheaf of \mathfrak{B} , then $\Gamma(U, \mathfrak{F}) = \{f \in \Gamma(U, \mathfrak{B}) | f_x \in \mathfrak{F}_x \text{ for all } x \in U\}$.

Lemma 2.3. *Let $G \neq \emptyset$ be an open subset of a complex manifold M . Let \mathfrak{B} be the sheaf of germs of bounded holomorphic functions on \bar{G} . Let $U \neq \emptyset$ be open in the space \bar{G} . Let \mathfrak{F} be a subsheaf of \mathfrak{B} . Suppose that $h \in \Gamma(U, \mathfrak{B})$ exists such that $\mathfrak{F}|U = h\mathfrak{B}|U$. Then for every $s \in \Gamma(U, \mathfrak{F})$, a function $g \in \Gamma(U, \mathfrak{B})$ exists such that $s = gh$. Observe: If V is any open subset of G whose closure \bar{V} is compact and contained in U , then g is bounded on $V \cap G$, i.e., $g|_{(V \cap G)} \in B_G(V)$.*

Proof. Pick $s \in \Gamma(U, \mathfrak{F})$. Then $s \in \Gamma(U, \mathfrak{B})$ with $s_x \in \mathfrak{F}_x$ for each $x \in U$. Hence for every $x \in U$ a germ $f(x) \in \mathfrak{B}_x$ exists such that $s_x = f(x) \cdot h_x$. An open neighborhood $W(x)$ of x in U exists such that $g^x \cdot h = s$ on $W(x) \cap G$ with $g^x \in B_G(W(x))$. If h is identically zero on some component of $W(x) \cap G$, then s is identically zero on this component; therefore it can be assumed that $g^x = 1$ on this component. Then $g^x = g^y$ on $W(x) \cap W(y) \cap G$ if this intersection is not empty. Hence one and only one holomorphic function g exists on $U \cap G$ such that $g|_{W(x) \cap G} = g^x$. Hence $g \cdot h = s$ on $U \cap G$. Let V be open in G such that \bar{V} is compact and contained in U . There, finitely many points x_1, \dots, x_p exist in U such that $\bar{V} \subseteq W(x_1) \cup \dots \cup W(x_p)$. Hence

$$\|g\|_{V \cap G} \leq \max_{v=1, \dots, p} \|g^{x_v}\|_{W(x_v) \cap G} < \infty.$$

Therefore $g|_{V \cap G} \in B_G(V)$. (Observe, that $\bar{V} \cap G$ may not be compact!)
O.E.D.

Now, Stout's result [6] can be formulated easily:

Theorem 2.4 (Stout). Let $G = G_1 \times \dots \times G_n$, where each $G_v = \{z \in \mathbb{C} | |z| < 1\}$ is the unit disc. Let \mathfrak{B} be the sheaf of germs of bounded holomorphic functions on \bar{G} . Then any principal ideal sheaf \mathfrak{F} in \mathfrak{B} is globally principal.

Proof. Finite, open coverings $\{U_i\}_{i \in I}$ and $\{V_i\}_{i \in I}$ of \bar{G} with $\bar{V} \subset U_i$ exist such that $\mathfrak{F}|_{U_i} = h_i \mathfrak{B}|_{U_i}$ with $h_i \in B_G(U_i)$. Here, \bar{V}_i is compact. Lemma 2.3 implies the existence of bounded holomorphic functions g_{ij} on $V_i \cap V_j \cap G$ if $V_i \cap V_j \neq \emptyset$ such that $h_i = g_{ij} h_j$ on $V_i \cap V_j \cap G$ where $g_{ij} = 1$ on every component of $V_i \cap V_j \cap G$ where $h_j \equiv 0$ and hence $h_i \equiv 0$. Then $g_{ji} = g_{ij}^{-1}$; especially, $g_{ij} \neq 0$ on $V_i \cap V_j \cap G$. Hence $\{h_i, V_i\}_{i \in I}$ is a Cousin II distribution in the sense of Stout [6]. Therefore, $h \in B_G(\bar{G})$ exists such that $h = u_i h_i$ and $h_i = v_i h$ on $V_i \cap G$ where u_i and v_i are bounded holomorphic functions on V_i . Clearly

$$\begin{aligned}\mathfrak{F}|_{V_i} &= h_i \mathfrak{B}|_{V_i} = h v_i \mathfrak{B}|_{V_i} \leq h \mathfrak{B}|_{V_i}, \\ h \mathfrak{B}|_{V_i} &= u_i h_i \mathfrak{B}|_{V_i} = u_i \mathfrak{F}|_{V_i} \leq \mathfrak{F}|_{V_i}.\end{aligned}$$

Hence $\mathfrak{F}|_{V_i} = h \mathfrak{B}|_{V_i}$. Since $\{V_i\}_{i \in I}$ is a covering, it follows that $\mathfrak{F} = h \mathfrak{B}$.
Q.E.D.

Theorem 2.5. Let $G = G_1 \times \cdots \times G_n \neq \emptyset$ be an open, bounded and connected subset of \mathbb{C}^n with $G_v \subset \mathbb{C}$ for $v = 1, \dots, n$. Let \mathfrak{B} be the sheaf of bounded holomorphic functions on \bar{G} . Let \mathfrak{F} be a globally principal ideal sheaf in \mathfrak{B} . Let $\mathcal{U} = \{U_i\}_{i \in I}$ be a finite, open covering of \bar{G} . Then

$$H^1(\bar{G}, \mathfrak{F}) = 0.$$

Proof. If \mathfrak{F} is the zero ideal, the statement is true. Suppose \mathfrak{F} is not the zero ideal. Take $h \in B_G(\bar{G})$ such that $\mathfrak{F} = h \mathfrak{B}$. Then h is not identically zero on G . Because G is connected, $h_x \neq 0$ for each $x \in \bar{G}$ and h_x is not a zero-divisor in \mathfrak{B}_x . If $g \in \mathfrak{F}_x$, then $g = u \cdot h_x$ with $u \in \mathfrak{B}_x$. Because h_x is not a zero divisor, u is unique. Define $u = \lambda(g)$. Obviously, $\lambda: \mathfrak{F} \rightarrow \mathfrak{B}$ is linear and bijective. It remains to be shown that λ is continuous. Pick $g_0 \in \mathfrak{F}_x$ and let W be an open neighborhood of $u_0 = \lambda(g_0) \in \mathfrak{B}_x$. An open neighborhood U of x in \bar{G} and $s \in B_G(U)$ and $t \in B_G(U)$ exist such that $s_x = u_0$ and $s_z \in W$ for all $z \in U$. Moreover, $t_x = g_0$ and $t_z \in \mathfrak{F}_z$ for all $z \in U$. Then $t_x = s_x h_x$. Hence $t = sh$ on $V \cap G$ for a neighborhood V of x in \bar{G} with $V \subseteq U$. Hence $t_z = s_z h_z$ for $z \in V$, which implies $\lambda(t_z) = s_z \in W$ for all $z \in V$. Now, $y = \{t_z | z \in V\}$ is an open neighborhood of $g_0 = t_x$ with $\lambda(y) \subseteq W$. Hence λ is continuous. Therefore $\lambda: \mathfrak{F} \rightarrow \mathfrak{B}$ is a sheaf isomorphism (for \mathfrak{B} -modules). Hence isomorphisms

$$\lambda^*: H^1(\mathcal{U}, \mathfrak{F}) \rightarrow H^1(\mathcal{U}, \mathfrak{B})$$

$$\lambda^*: H^1(\bar{G}, \mathfrak{F}) \rightarrow H^1(\bar{G}, \mathfrak{B})$$

are induced. Now, Theorem 2.1 and Theorem 2.2 complete the proof. Q.E.D.

Let $G \neq \emptyset$ be an open subset of the complex manifold M . Let \mathfrak{B} be the sheaf of germs of bounded holomorphic functions on M . Let \mathfrak{I} be a principal ideal sheaf in \mathfrak{B} . Let \mathfrak{Q} be the quotient sheaf. This defines the exact sequence.

$$0 \rightarrow \mathfrak{I} \rightarrow \mathfrak{B} \xrightarrow{\mu} \mathfrak{Q} \rightarrow 0$$

where μ is the residual map. Observe that \mathfrak{I} , \mathfrak{B} and \mathfrak{Q} are sheaves on \bar{G} , not only on G . Then

$$\text{supp } \mathfrak{Q} = \{x \in \bar{G} \mid \mathfrak{Q}_x \neq 0\} = \{x \in \bar{G} \mid \mathfrak{I}_x \neq \mathfrak{B}_x\}.$$

The intersection $S(\mathfrak{I}) = G \cap \text{supp } \mathfrak{Q}$ is analytic. If $\mathfrak{I} \neq \mathfrak{B}$, and if $\mathfrak{I}_x \neq 0$ for at least one point in each component of G , then $S(\mathfrak{I})$ has pure codimension 1. Observe that $(\text{supp } \mathfrak{Q}, \mathfrak{Q})$ is a ringed space. \mathfrak{Q}_x may have zero divisors. For each open subset U of the space \bar{G} with $V = U \cap \text{supp } \mathfrak{Q} \neq \emptyset$, $\Gamma(V, \mathfrak{Q}) = \Gamma(U, \mathfrak{Q})$ is a ring. Its elements are, so to speak, the "holomorphic functions" on U in $(\text{supp } \mathfrak{Q}, \mathfrak{Q})$. However, in general, this ring cannot be identified with a ring of functions on V .

If N is any analytic subset of G , and if $V \neq \emptyset$ is open in the space N , let $A(V; N)$ be the ring of holomorphic functions on V in the sense of Serre. If U is open in the space \bar{G} and if $V = U \cap \text{supp } \mathfrak{Q} \neq \emptyset$, then a ring homomorphism

$$v: \Gamma(V, \mathfrak{Q}) \rightarrow A(S(\mathfrak{I}) \cap V, S(\mathfrak{I}))$$

shall be defined: Take $s \in \Gamma(V, \mathfrak{Q})$. For every $x \in V$ an open neighborhood W_x of x in the space \bar{G} and $f^x \in B_G(W_x)$ exist such that $\mu(f_z^x) = s(z)$ for all $z \in W_x$, where $s(z) = 0$ for $z \in U - V$. If $W_x \cap W_y \neq \emptyset$, then $f_z^x - f_z^y \in \mathfrak{I}_z$ for all $z \in W_x \cap W_y$. Hence the function

$$v(s): S(\mathfrak{I}) \cap V \rightarrow \mathbb{C}$$

is well defined by $v(s)(z) = f^y(z)$ if $z \in W_x \cap S(\mathfrak{I})$. Obviously, $v(s) \in A(S(\mathfrak{I}) \cap V, S(\mathfrak{I}))$. Obviously, v is a ring-homomorphism, but, in general, v is not injective or surjective. $v(s)(z)$ is said to be the *value of s at z* . If $x \in V$, then $f^x|_{W_x \cap S(\mathfrak{I})} = v(s)|_{W_x \cap S(\mathfrak{I})}$. Hence, for each point $x \in V$ a neighborhood $W = W_x$ exists such that $v(s)$ is bounded on $W_x \cap S(\mathfrak{I})$. Hence, if \bar{G} is compact, and $s \in \Gamma(\bar{G}, \mathfrak{Q})$, then $v(s)$ is bounded on $S(\mathfrak{I})$, because $\text{supp } \mathfrak{Q}$ is compact and can be covered by finitely many neighborhoods W_x . In general, not every bounded holomorphic function on $S(\mathfrak{I})$ can be obtained this way. (See the example of Alexander at the end of this paper.)

A holomorphic function $f \in A(G, S(\mathfrak{F}))$ is said to be *strictly bounded* for \mathfrak{F} if there exists an open covering $\mathfrak{U} = \{U_i\}_{i \in I}$ of the space \bar{G} and a family $\{f_i\}_{i \in I}$ of functions $f_i \in B_G(U_i)$ such that:

- 1) If $(i, j) \in I(1)$ and if $z \in U_i \cap U_j$, then $(f_i - f_j)_z \in \mathfrak{F}_z$.
- 2) If $i \in I$ with $U_i \cap S(\mathfrak{F}) \neq \emptyset$ then $f_i|_{U_i \cap S(\mathfrak{F})} = f|_{U_i \cap S(\mathfrak{F})}$.

If \bar{G} is compact, finitely many U_i cover \bar{G} and f is bounded.

Lemma 2.6. *The function $f \in A(G, S(\mathfrak{F}))$ is strictly bounded for \mathfrak{F} , if and only if $f = v(s)$ for some $s \in \Gamma(\text{supp } \mathfrak{Q}, \mathfrak{Q})$.*

Proof. a) If f is strictly bounded, define s by $s(z) = \mu(f_{iz})$ for $z \in U_i$. Because of 1), $s(z)$ does not depend on i . Hence s is well defined: $s: \bar{G} \rightarrow \mathfrak{Q}$. Because the germ $f_{iz} \in \mathfrak{B}$ depends continuously on $z \in U_i$, the section s is continuous. By definition and because of 2), $v(s) = f$.

b) Suppose that $f = v(s)$ with $s \in \Gamma(\bar{G}, \mathfrak{Q}) = \Gamma(\text{supp } \mathfrak{Q}, \mathfrak{Q})$. For every $x \in \bar{G}$, an open neighborhood W_x of x in the space \bar{G} and $f^x \in B_G(W_x)$ exists such that $\mu(f_z^x) = s(z)$ for all $z \in W_x$. If $W_x \cap W_y \neq \emptyset$ then $f_z^x - f_z^y \in \mathfrak{F}_z$ for all $z \in W_x \cap W_y$. Moreover,

$$f|_{W_x \cap S(\mathfrak{F})} = v(s)|_{W_x \cap S(\mathfrak{F})} = f^x|_{W_x \cap S(\mathfrak{F})}. \quad \text{Q.E.D.}$$

Lemma 2.7. *The holomorphic function $f \in A(G, S(\mathfrak{F}))$ is strictly bounded for \mathfrak{F} , if and only if an open covering $\mathfrak{U} = \{U_i\}_{i \in I}$ of the space \bar{G} , a family $\{f_i\}_{i \in I}$ with $f_i \in B_G(U_i)$, and functions $h_{ij} \in B_G(U_i \cap U_j)$ for each $(i, j) \in I(1)$ exist such that:*

1. If $i \in I$ and if $U_i \cap S(\mathfrak{F}) \neq \emptyset$, then $f|_{U_i \cap S(\mathfrak{F})} = f_i|_{U_i \cap S(\mathfrak{F})}$.
2. If $(i, j) \in I(1)$, then $f_i - f_j = a_{ij}h_{ij}$ on $U_i \cap U_j \cap G$.
3. If $(i, j) \in I(1)$, then $\mathfrak{F}|_{U_i \cap U_j} = h_{ij}\mathfrak{B}|_{U_i \cap U_j}$.

Proof. a) Obviously, if these conditions are satisfied, then f is strictly bounded.

b) Let f be strictly bounded. An open covering $\mathfrak{U} = \{U_i\}_{i \in I}$ of the space \bar{G} and family $\{f_i\}_{i \in I}$ of functions $f_i \in B_G(U_i)$ exist such that the conditions of the definition of "strictly bounded" hold. These remain true for any refinement. Hence it can be assumed that $\mathfrak{F}|_{U_i}$ is generated by an element of $B_G(U_i)$ such that there exists an open covering $\mathfrak{B} = \{V_i\}_{i \in I}$ of \bar{G} with \bar{V}_i compact and contained in U_i . Now, if $(i, j) \in I(1)$, an element $h_{ij} \in B_G(U_i \cap U_j)$ exists such that $\mathfrak{F}|_{U_i \cap U_j} = h_{ij}\mathfrak{B}|_{U_i \cap U_j}$. Then $(f_i - f_j)_z \in \mathfrak{F}_z$ for all $z \in U_i \cap U_j$. Lemma 2.3 gives $a_{ij} \in B_G(V_i \cap V_j)$ such that $f_i - f_j = a_{ij}h_{ij}$ on $V_i \cap V_j \cap G$. Hence the conditions 1-3 of the Lemma are satisfied for the covering \mathfrak{B} , the family $\{f_i|_{V_i \cap G}\}_{i \in I}$ and the functions $h_{ij}|_{V_i \cap V_j \cap G}$ and a_{ij} (observe that it was necessary to use Lemma 2.3). Q.E.D.

Now, the main result can be proved:

Theorem 2.8. *Let $G = G_1 \times \cdots \times G_n \neq \emptyset$ be an open, bounded and connected subset of \mathbb{C}^n with $G_v \subset \mathbb{C}$ for $v = 1, \dots, n$. Let \mathfrak{B} be the sheaf of bounded holomorphic functions on \bar{G} . Let \mathfrak{I} be a globally principal ideal sheaf in \mathfrak{B} with $\mathfrak{I} \neq \mathfrak{B}$. Let f be a strictly bounded holomorphic function for \mathfrak{I} on $S(\mathfrak{I})$. Then a bounded holomorphic function F on G exists such that*

$$F|_{S(\mathfrak{I})} = f.$$

Proof. The exact sequence $0 \rightarrow \mathfrak{I} \rightarrow \mathfrak{B} \rightarrow \mathfrak{Q} \rightarrow 0$ induces an exact sequence

$$0 \rightarrow \Gamma(\bar{G}, \mathfrak{I}) \rightarrow \Gamma(\bar{G}, \mathfrak{B}) \xrightarrow{\tilde{\mu}} \Gamma(\bar{G}, \mathfrak{Q}) \rightarrow H^1(\bar{G}, \mathfrak{I}) = 0.$$

Hence $\tilde{\mu}$ is surjective. Take $s \in \Gamma(\bar{G}, \mathfrak{Q})$ such that $v(s) = f$. A bounded holomorphic function $F \in B_G(\bar{G}) = \Gamma(\bar{G}, \mathfrak{B})$ exists such that $\tilde{\mu}(F) = s$. Hence $\mu(F_z) = s(z)$ for each $z \in \bar{G}$. Hence $f = v(s) = F|_{S(\mathfrak{I})}$ by the construction of $v(s)$ using $W_x = \bar{G}$ and $f^x = F$ for all $x \in \bar{G}$. Q.E.D.

Theorem 2.8 and Stout's Theorem 2.4 imply:

Theorem 2.9. *Let $G = G_1 \times \cdots \times G_n$ where each $G_v = \{z \in \mathbb{C} \mid |z| < 1\}$ is the unit disc. Let \mathfrak{B} be the sheaf of germs of bounded holomorphic functions on \bar{G} and \mathfrak{I} be a principal ideal sheaf in \mathfrak{B} with $\mathfrak{I} \neq \mathfrak{B}$. Let f be a strictly bounded holomorphic function for \mathfrak{I} on $S(\mathfrak{I})$. Then a bounded holomorphic function F exists on G such that $F|_{S(\mathfrak{I})} = f$.*

Define $G = \{(z, w) \in \mathbb{C}^2 \mid |z| < 1 \text{ and } |w| < 1\}$. Define $h(z, w) = (w - \frac{1}{2})(zw - \frac{1}{2})$. Define $\mathfrak{I} = h\mathfrak{B}$ on G . Define

$$S_1 = \{(z, \frac{1}{2}) \mid z \in \mathbb{C} \text{ with } |z| < 1\},$$

$$S_2 = \{(z, w) \in G \mid zw = \frac{1}{2}\}.$$

Then $S(\mathfrak{I}) = S_1 \cup S_2$. Define $f: S(\mathfrak{I}) \rightarrow \mathbb{C}$ by $f|_{S_1} = 0$ and $f|_{S_2} = 1$. Then f is bounded and holomorphic. Alexander [1] has shown that f is not the restriction of a bounded holomorphic function on G . Hence f is not strictly bounded!

NOTES

1. In this research, the first author was partially supported by the Air Force under Grant AF 49(638)1345. The second author was partially supported by the National Science Foundation under Grant NSF G 72 65, while both authors were visitors at Stanford University.

2. See also Siu [5].

REFERENCES

- [1] ALEXANDER, H., Extending bounded holomorphic functions from certain subvarieties of a polydisc, *Pacific J. Math.* **29**, (1969), 485-490.
- [2] GRAUERT, H., AND I. LIEB, Das Ramirezsche Integral und die Lösung der Gleichung $\bar{\partial}f = a$ im Bereich der beschränkten Formen (Complex Analysis (1969)), *Rice Univ. Studies* **56**, No. 2 (1970), 29-50.
- [3] HÖRMANDER, L., An Introduction to Complex Analysis in Several Variables, D. Van Nostrand Co., Princeton, N. J. (1966).
- [4] RAMIREZ DE ARELLANO, E., Ein Divisionsproblem in der komplexen Analysis mit einer Anwendung auf Randintegraldarstellungen, *Math. Ann.* **184** (1970), 172-187.
- [5] SIU, Y.-T., Sheaf cohomology with bounds and bounded holomorphic functions, *Proc. Amer. Math. Soc.* **21** (1969), 226-229.
- [6] STOUT, E. L., The second Cousin problem with bounded data, *Pacific J. Math.* **26** (1968), 379-387.

UNIVERSITY OF PISA
UNIVERSITY OF NOTRE DAME